

Chapter 6: Searching (Cont.)

Searching by Elimination
– Program Refinement –

Searching by Elimination

Given:

- a finite set W
- a boolean function S on W , such that $S.w$ holds for some $w \in W$

derive a program with the following post-condition.

$$S.x$$

Note: if S is identified with $\{x \in W \mid S.x\}$, then the post-condition can be written as

$$R: S \cap \{x\} \neq \emptyset.$$

What is a suitable invariant?

Generalization of the post-condition:

$$R : S \cap \underline{\{x\}} \neq \emptyset.$$

gives as invariant

$$P : S \cap \underline{V} \neq \emptyset \wedge \underline{V} \subseteq W$$

which is established by $V := W$.

This leads to the following program scheme:

$\{S \cap W \neq \emptyset\}$

$V := W;$

{invariant: $S \cap V \neq \emptyset \wedge V \subseteq W$, bound: $|V|$ }

do $|V| \neq 1 \rightarrow$

 decrease $|V|$ under invariance of P

od;

$x :=$ the unique element of V

Searching by elimination:

```

{ $S \cap W \neq \emptyset$ }
 $V := W$ ;
{invariant:  $S \cap V \neq \emptyset \wedge V \subseteq W$ , bound:  $|V|$ }
do  $|V| \neq 1 \rightarrow$ 
    choose  $a$  and  $b$  in  $|V|$  such that  $a \neq b$ 
    { $a \in V \wedge b \in V \wedge a \neq b \wedge S \cap V \neq \emptyset$ }
    if  $B_0 \rightarrow V := \underline{V \setminus \{a\}}$ 
        []  $B_1 \rightarrow V := \underline{V \setminus \{b\}}$ 
    fi
od;
 $x :=$  the unique element of  $V$ 

```

What are B_0 and B_1 ?

B_0 and B_1 should be the conditions keeping invariants, i.e.,

$$B_0 \Rightarrow (S \cap V \neq \emptyset \Rightarrow S \cap (V \setminus \{a\}) \neq \emptyset)$$

$$B_1 \Rightarrow (S \cap V \neq \emptyset \Rightarrow S \cap (V \setminus \{b\}) \neq \emptyset)$$

How to calculate out B_0 and B_1 ?

From the calculation

$$\begin{aligned}
 & S \cap V \neq \emptyset \Rightarrow S \cap (V \setminus \{a\}) \neq \emptyset \\
 \equiv & \{ a \in V \} \\
 & S.a \vee (S \cap (V \setminus \{a\})) \neq \emptyset \Rightarrow S \cap (V \setminus \{a\}) \neq \emptyset \\
 \equiv & \{ \text{predicate calculus} \} \\
 & S.a \Rightarrow S \cap (V \setminus \{a\}) \neq \emptyset \\
 \Leftarrow & \{ b \in V \setminus \{a\} \} \\
 & S.a \Rightarrow S.b \\
 \equiv & \{ \text{prediate calculus} \} \\
 & \neg S.a \vee S.b
 \end{aligned}$$

we may have

$$B_0 = \neg S.a \vee S.b.$$

And similarly we may have

$$B_1 = \neg S.b \vee S.a.$$

So we obtain the program:

```

{ $S \cap W \neq \emptyset$ }
 $V := W$ ;
{invariant:  $S \cap V \neq \emptyset \wedge V \subseteq W$ , bound:  $|V|$ }
do  $|V| \neq 1 \rightarrow$ 
    choose  $a$  and  $b$  in  $|V|$  such that  $a \neq b$ 
    { $a \in V \wedge b \in V \wedge a \neq b \wedge S \cap V \neq \emptyset$ }
    if  $\neg S.a \vee S.b \rightarrow V := \underline{V \setminus \{a\}}$ 
    []  $\neg S.b \vee S.a \rightarrow V := \underline{V \setminus \{b\}}$ 
    fi
od;
 $x :=$  the unique element of  $V$ 

```

Refinement

A special case when $W = [0..N]$. We may choose V can be represented by two integers a and b as $[a..b]$, and the program becomes

```
{(∃i : 0 ≤ i ≤ N : S.i)}  
a, b := 0, N;  
do a ≠ b →  
    if ¬S.a ∨ S.b → a := a + 1  
    [] ¬S.b ∨ S.a → b := b - 1  
    fi  
od;  
x := a;
```

Application 1:

Derive a program that satisfies

```
[[  
  con  $N : int \{N \geq 0\}; b : \mathbf{array} [0..N] \mathbf{of} int;$   
  var  $x : int;$   
  max location  
   $\{0 \leq x \leq N \wedge f.x = (\mathbf{max} i : 0 \leq i \leq N : f.i)\}$   
  ]]  
   $\{S.x\}$ 
```

How to make use of “Searching by Elimination” to solve this problem?

The post-condition can be rewritten as

$$R : 0 \leq x \leq N \wedge (\forall i : 0 \leq i \leq N : f.i \leq f.x)$$

and we can define S as

$$S.x \equiv (\forall i : 0 \leq i \leq N : f.i \leq f.x)$$

What is a sufficient condition for $\neg S.a \vee S.b$?

Since

$$\begin{aligned}
 & \neg S.a \vee S.b \\
 \equiv & \quad \{ \text{predicate calculus} \} \\
 & S.a \Rightarrow S.b \\
 \equiv & \quad \{ \text{definition of } S \} \\
 & (\forall i : 0 \leq i \leq N : f.i \leq f.a) \Rightarrow (\forall i : 0 \leq i \leq N : f.i \leq f.b) \\
 \Leftarrow & \quad \{ \text{transitivity of } \leq \} \\
 & f.a \leq f.b
 \end{aligned}$$

we have

$$f.a \leq f.b \Rightarrow \neg S.a \vee S.b$$

Similarly, we can derive

$$f.b \leq f.a \Rightarrow \neg S.b \vee S.a$$

Our solution:

```
var  $a, b : int;$   
 $a, b := 0, N;$   
do  $a \neq b \rightarrow$   
    if  $f.a \leq f.b \rightarrow a := a + 1$   
     $\square$   $f.b \leq f.a \rightarrow b := b - 1$   
    fi  
od;  
 $x := a;$ 
```

Application 2: The Celebrity Problem

Design a program to compute a *celebrity* among $N + 1$ persons. A person is a celebrity if he is known by everyone but does not know anyone.

```

[[
  con  $N : int \{N \geq 0\}$ ;  $k : array [0..N] \times [0..N] \text{ of } bool$ ;
   $\{(\exists i : 0 \leq i \leq N : (\forall j : j \neq i : k.j.i \wedge \neg k.i.j))\}$ 
  var  $x : int$ ;
  celebrity
   $\{0 \leq x \leq N \wedge (\forall j : j \neq x : k.j.x \wedge \neg k.x.j)\}$ 
]]

```

Here $k.i.j$ denotes i knows j .

We could consider the set W as $[0..N]$. What is S ?

We choose

$$S.x \equiv (\forall j : j \neq x : k.j.x \wedge \neg k.x.j)$$

We then derive

$$\begin{aligned} & \neg S.a \vee S.b \\ \Leftarrow & \quad \{ \text{predicate calculus} \} \\ & \neg S.a \\ \equiv & \quad \{ \text{definition of } S \} \\ & \neg(\forall j : j \neq a : k.j.a \wedge \neg k.a.j) \\ \equiv & \quad \{ \text{De Morgan} \} \\ & (\exists j : j \neq a : \neg k.j.a \vee k.a.j) \\ \Leftarrow & \quad \{ b \neq a \} \\ & \neg k.b.a \vee k.a.b \end{aligned}$$

We thus obtain the following program:

```
{(∃i : 0 ≤ i ≤ N : S.i)}  
a, b := 0, N;  
do a ≠ b →  
    if ¬k.b.a ∨ k.a.b → a := a + 1  
    [] ¬k.a.b ∨ k.b.a → b := b - 1  
    fi  
od;  
x := a;
```

Chapter 7: Segment Problems

This chapter is to show

- how problems may be solved;
- what decisions are made in the derivations;
- which properties play a specific role.

Longest Segment Problems

Let $N \geq 0$ and let $X[0..N)$ be an integer array. Find the longest subsegment $[p..q)$ of $[0..N)$ that satisfies a certain predicate like

- all elements are zero: $(\forall i : p \leq i < q : X.i = 0)$
- the segment is left-minimal: $(\forall i : p \leq i < q : X.p \leq X.i)$
- the segment contains at most 10 zeros: $(\#i : p \leq i < q : X.i = 0) \leq 10$
- all values are different: $(\forall i, j : p \leq i < j < q : X.i \neq X.j)$

All Zeros

Determine the length of a longest segment of $X[0..N)$ that contains zeros only.

```
[[  
  con  $N : int \{N \geq 0\}; X : \mathbf{array} [0..N) \mathbf{of} int;$   
  var  $r : int;$   
  all zeros  
   $\{r = (\mathbf{max} p, q : 0 \leq p \leq q \leq N \wedge (\forall i : p \leq i < q : X.i = 0) : q - p)\}$   
  ]]
```

The post-condition is:

$$R : r = (\mathbf{max} p, q : 0 \leq p \leq q \leq N \wedge A.p.q : q - p)$$

where

$$A.p.q = (\forall i : p \leq i < q : X.i = 0)$$

What properties does A have?

For

$$A.p.q = (\forall i : p \leq i < q : X.i = 0)$$

we have

- A holds for empty segments: $A.n.n$
- A is prefix-closed: $A.p.q \Rightarrow (\forall i : p \leq i \leq q : A.p.i)$
- A is postfix-closed: $A.p.q \Rightarrow (\forall i : p \leq i \leq q : A.i.q)$

Our invariants are from R by replacing constant N by variable n :

$$P_0 : r = (\mathbf{max} p, q : 0 \leq p \leq q \leq n \wedge A.p.q : q - p)$$

$$P_1 : 0 \leq n < N$$

which is established by $n, r := 0, 0$.

What if $n := n + 1$?

$$\begin{aligned}
& (\mathbf{max} \ p, q : 0 \leq p \leq q \leq n + 1 \wedge A.p.q : q - p) \\
= & \quad \{ \text{split off } q = n + 1 \} \\
& (\mathbf{max} \ p, q : 0 \leq p \leq q \leq n \wedge A.p.q : q - p) \mathbf{max} \\
& \quad (\mathbf{max} \ p, q : 0 \leq p \leq n + 1 \wedge A.p.(n + 1) : n + 1 - p) \\
= & \quad \{ P_0 \} \\
& r \mathbf{max} (\mathbf{max} \ p : 0 \leq p \leq n + 1 \wedge A.p.(n + 1) : n + 1 - p) \\
= & \quad \{ + \text{ distributes over } \mathbf{max} \} \\
& r \mathbf{max} (n + 1 + (\mathbf{max} \ p : 0 \leq p \leq n + 1 \wedge A.p.(n + 1) : - p)) \\
= & \quad \{ \text{property of } \mathbf{max} \text{ and } \mathbf{min} \} \\
& r \mathbf{max} (n + 1 - (\mathbf{min} \ p : 0 \leq p \leq n + 1 \wedge A.p.(n + 1) : p)) \\
= & \quad \{ \text{invariant strengthening: } Q : s = (\mathbf{min} \ p : 0 \leq p \leq n \wedge A.p.n : p) \} \\
& r \mathbf{max} (n + 1 - s)
\end{aligned}$$

We thus obtain a program of the following form.

```
 $n, r, s := 0, 0, 0; \{ \text{invariant: } P_0 \wedge P_1 \wedge Q, \text{ bound: } N - n \}$   
do  $n \neq N \rightarrow$   
    establish  $Q(n := n + 1)$   
     $r := r \mathbf{max} (n + 1 - s);$   
     $n := n + 1$   
od
```

How to solve the subproblem establishing $Q(n := n + 1)$:

$$Q : s = (\mathbf{min} p : 0 \leq p \leq n \wedge A.p.n : p)$$

We may remove **min** to the conjunction of the following predicates:

$$Q_0 : 0 \leq s \leq n$$

$$Q_1 : A.s.n$$

$$Q_2 : (\forall p : 0 \leq p < s : \neg A.p.n)$$

Lemma. If A is prefix-closed, then

$$Q_0 \wedge Q_2 \wedge A.s.(n + 1) \Rightarrow Q(n := n + 1)$$

From

$$Q_0 \wedge Q_2 \wedge A.s.(n + 1) \Rightarrow Q(n := n + 1)$$

we can establish $Q(n := n + 1)$ by considering

- $Q_0 \wedge Q_2$ as invariant,
- $\neg A.s.(n + 1)$ as guard, and
- $n + 1 - s$ as bound function.

Theorem

```

[[
{A holds for empty segment and prefix-closed}
var  $n, s : int$ ;
 $n, r, s := 0, 0, 0$ ;
do  $n \neq N \rightarrow$ 
    do  $\neg A.s.(n + 1) \rightarrow s := s + 1$  od;
     $r := r \mathbf{max} (n + 1 - s)$ ;
     $n := n + 1$ 
od
{ $r = (\mathbf{max} p, q : 0 \leq p \leq q \leq N \wedge A.p.q : q - p)$ }
]]

```

What is its time complexity?

Add the variable t for counting the steps.

```

[[
var  $n, s, t : int$ ;
 $n, r, s, t := 0, 0, 0, 0$ ;
{invariant :  $Q : s = (\mathbf{min} p : 0 \leq p \leq n \wedge A.p.n : p)$ }
do  $n \neq N \rightarrow$ 
    do  $\neg A.s.(n + 1) \rightarrow s := s + 1; t := t + 1$  od;
     $r := r \mathbf{max} (n + 1 - s)$ ;
     $n := n + 1; t := t + 1$ 
od
]]

```

It is not difficult to see that $t = s + n \leq 2N$. So if $A.s.(n + 1)$ can be computed in constant time, then the above program is linear.

For the all-zeros problem, how $A.s.(n + 1)$ can be computed in constant time?

from

$$A.p.q = (\forall i : p \leq i < q : X.i = 0)?$$

Point: try to make use the the invariant:

$$Q : s = (\mathbf{min} p : 0 \leq p \leq n \wedge A.p.n : p)$$

Let us investigate the effect of $A.s.(n + 1)$:

$$\begin{aligned}
 & A.s.(n + 1) \\
 \equiv & \quad \{ \text{definition of } A \} \\
 & (\forall i : s \leq i < n + 1 : X.i = 0) \\
 \equiv & \quad \{ \text{split off } i = n, \underline{s \leq n} \} \\
 & (\forall i : s \leq i < n : X.i = 0) \wedge X.n = 0 \\
 \equiv & \quad \{ \text{definition of } A \} \\
 & A.s.n \wedge X.n = 0 \\
 \equiv & \quad \{ \text{by the invariant } Q \} \\
 & X.n = 0
 \end{aligned}$$

The final program:

```
[[  
  var  $n, s : int$ ;  
   $n, r, s := 0, 0, 0$ ;  
  do  $n \neq N \rightarrow$   
    do  $X.n \neq 0 \wedge s \leq n \rightarrow s := s + 1$  od;  
     $r := r \max (n + 1 - s)$ ;  
     $n := n + 1$   
  od  
]]
```

Exercises

[Problem 8-1] The Starting Pit Location Problem

Given are $N + 1$ pits located along a circular racetrack. The pits are numbered clockwise from 0 up to and including N . At pit i , there are $p.i$ gallons of petrol available. To race from pit i to its clockwise neighbor one needs $q.i$ gallons of petrol. One is asked to design a linear algorithm to determine a pit from which it is possible to race a complete lap, starting with an empty fuel tank. To guarantee the existence of such a starting pit, we assume that

$$(\sum_{i : 0 \leq i \leq N} p.i) = (\sum_{i : 0 \leq i \leq N} q.i).$$

[Problem 8-2]

Derive an $O(N)$ solution to the following problem.

[[

con $N : int \{N \geq 0\}; X : \text{array } [0..N) \text{ of } int;$

var $r : int;$

all equal

$\{r = (\mathbf{max} \ p, q : 0 \leq p \leq q \leq N \wedge (\forall i, j : p \leq i \leq j < q : X.i = X.j) : q - p)\}$

]]

About the Final Report

- Solve 8 problems freely selected from the exercises in the lecture notes.
- Submit your report to my post-box in the first floor of Engineering Building 6 no later than 17pm, June 19 (Thursday), 2008. Never forget writing your name, student identification number, and your department.
- Note you will be asked to present one or two of your solutions in class.